

BRONKOXT

Bromhexine MUCOLYTIC EXPECTORANT ORAL SOLUTION

PHARMACEUTICAL FORM: ORAL SOLUTION

COMPOSITION:

Each 100 ml contains:

Bromhexine 0.6 g
Excipients q.s. 100 mL

SPECIES:

- Feline
- Canine

DOSAGE

DOSAGE		
SPECIE	REFERENCE DOSE	PRACTICAL DOSE
 Canine	2 mg/kg	3,3 mL/10 kg/q12h for 7-10 days
 Feline	1 mg/kg	0,5 mL/3kg/q 24h for 7-10 days
Route of administration: Oral		

INDICATIONS AND USE:

BRONKOXT is a direct-acting mucolytic-expectorant, bromhexine, stimulates and facilitates the expulsion of tracheo-bronchial secretions by decreasing the viscosity of secretions, allowing rapid evacuation. Is indicated for the treatment of all respiratory tract conditions, infectious or non-infectious, acute or chronic with mucus discharge.

ROUTE OF ADMINISTRATION ORAL

PHARMACOKINETICS

It is metabolized at the liver level, it is transformed into inactive metabolites that are eliminated through the feces and urine. In dogs, after oral administration,

it experiences a rapid absorption, reaching the plasma maximum within 0.5-2 hours of administration, although the bioavailability is very low (less than 10%) because first-pass metabolism is very efficient. Elimination is greater by fecal route in canines than in other species, where urinary predominates.

MECHANISM OF ACTION:

Regulates the production of bronchial mucus, often increase its net quantity and modifying the action of the secretory cells. Reduce the viscosity of mucus and increases tracheobronchial ciliary activity.

PRECAUTION, WARNINGS AND CONTRAINDICATIONS

- Use fresh solutions.
- Do not administer in animals with gastric ulcers.

DRUG INTERACTIONS

- Risks of mucus stasis due to inhibition of the cough reflex with: antitussives (anticholinergics, antihistamines, etc.).
- Antagonized effect by inhibiting bronchial secretion with: anticholinergics, H1 antihistamines, tricyclic antidepressants, neuroleptics.

SIDE EFFECTS

It could occur in animals with liver failure and severe kidney

ANTIDOTE

- No information

STORAGE

Store in a cool and dry place, protect from light at a temperature between 15°C to 30°C.

Keep out of the reach of children and pets.

SALE

Veterinary prescription only.

Ecuador Reg. N°: 6E-12472-AGROCALIDAD
Kingdom of Bahrain Distributor in GCC
Arsan for veterinary services
Info@arsan.vet
+973 37773682

Elaborated and distributed by: **FARBIOPHARMA S.A.**
Address: Guayas Street N° E3-296 and Pichincha Ave.
Location: Lomas de la Concepción, way to Pintag.
Phone: (593) 2 5240501 - 2794046 - 2794031
+593 99 968 4047 / +593 99 184 6785
CUSTOMER SERVICE: 1800327246
www.farbiopharma.com / Quito - Ecuador

MADE IN ECUADOR